

XIAFLEX® (collagenase clostridium histolyticum) Coding Scenario 2: Procedure Modifier Required



HEALTH INSURANCE CLAIM FORM

Medicare Part B Claims

APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12

<input type="checkbox"/> PICA										PICA <input type="checkbox"/>																																		
1. MEDICARE <input type="checkbox"/> (Medicare#)					MEDICAID <input type="checkbox"/> (Medicaid#)					TRICARE <input type="checkbox"/> (ID#/DoD#)					CHAMPVA <input type="checkbox"/> (Member ID#)					GROUP HEALTH PLAN <input type="checkbox"/> (ID#)					FECA BLK LUNG <input type="checkbox"/> (ID#)					OTHER <input type="checkbox"/> (ID#)					1a. INSURED'S I.D. NUMBER (For Program in Item 1) 123-45-6789A									
2. PATIENT'S NAME (Last Name, First Name, Middle Initial) Doe, John Q.															3. PATIENT'S BIRTH DATE MM DD YY 10 19 1935										SEX <input checked="" type="checkbox"/> M <input type="checkbox"/> F					4. INSURED'S NAME (Last Name, First Name, Middle Initial) Same														
5. PATIENT'S ADDRESS (No., Street) 1212 Main St.															6. PATIENT RELATIONSHIP TO INSURED Self <input checked="" type="checkbox"/> Spouse <input type="checkbox"/> Child <input type="checkbox"/> Other <input type="checkbox"/>										7. INSURED'S ADDRESS (No., Street) Same																			
CITY Any City					STATE XX					8. RESERVED FOR NUCC USE										CITY					STATE																			
ZIP CODE XXXXX					TELEPHONE (Include Area Code)										ZIP CODE					TELEPHONE (Include Area Code)																								
9. OTHER INSURANCE										10. RESERVED FOR NUCC USE										11. RESERVED FOR NUCC USE																								
a. OTHER INSURANCE AARP										b. RESERVED FOR NUCC USE										c. RESERVED FOR NUCC USE																								
d. INSURANCE										12. PATIENT'S AUTHORITY TO PROCESS BELOW										13. RESERVED FOR NUCC USE																								
12. PATIENT'S AUTHORITY TO PROCESS BELOW SIGNED _____										14. DATE OF SIGNATURE MM DD YY										17. NAME OF SUPPLIER																								
14. DATE OF SIGNATURE MM DD YY										17. NAME OF SUPPLIER NPI										FROM TO																								
19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)										20. OUTSIDE LAB? <input type="checkbox"/> YES <input type="checkbox"/> NO \$ CHARGES										22. RESUBMISSION CODE ORIGINAL REF. NO.																								
21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to service line below (24E) ICD Ind.										A. M72.0										23. PRIOR AUTHORIZATION NUMBER																								
24. A. DATE(S) OF SERVICE From To B. PLACE OF SERVICE C. EMG D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances) CPT/HCPCS MODIFIER E. DIAGNOSIS POINTER F. \$ CHARGES G. DAYS OR UNITS H. EPSON Family Plan I. ID. QUAL. J. RENDERING PROVIDER ID. #										10 03 21 10 03 21 11 J0775 A XXX XX 116 NPI										10 03 21 10 03 21 11 J0775 JW A XXX XX 64 NPI																								
10 03 21 10 03 21 11 20527 A XXX XX 1 NPI										10 03 21 10 03 21 11 20527 A XXX XX 1 NPI										10 05 21 10 05 21 11 26341 A XXX XX 1 NPI																								
10 05 21 10 05 21 11 26341 A XXX XX 1 NPI										25. FEDERAL TAX I.D. NUMBER SSN EIN										26. PATIENT'S ACCOUNT NO.																								
123456789										27. ACCEPT ASSIGNMENT? (For gov. claims, see back) <input type="checkbox"/> YES <input checked="" type="checkbox"/> NO										28. TOTAL CHARGE \$ xxx.xx										29. AMOUNT PAID \$ xxx.xx														
31. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREES OR CREDENTIALS (I certify that the statements on the reverse apply to this bill and are made a part thereof.)										32. SERVICE FACILITY LOCATION INFORMATION										30. Rsvd for NUCC Use																								
SIGNED _____ DATE _____										a. NPI b.										00 (02-12)																								

SAMPLE 1500 FORM TREATMENT BILLING WHEN TREATING 2 CORDS/JOINTS

Based on prior policy, this sample represents how your Medicare Administrative Contractor (MAC) is likely to require completion of claim forms for J0775, and the CPT® codes 20527 and 26341. This sample form is not intended to replace or modify your MAC's policy, and use of this form does not guarantee payment or take the place of professional coding advice. Coding is part of the clinical decision, and each provider is responsible for selecting the billing codes that most accurately describe the services provided, and for adhering to all payor guidance. Information is subject to change. Please refer to the policy on the MAC's website. This sample claim form does not represent any clinical or treatment recommendation.

Use JW modifier to indicate how much drug is wasted from single-use vials.

These amounts represent 58 units per cord/joint (total of 116 units) and wastage of 32 units per cord/joint (total of 64 units) and are provided as an example of the recommended dose and wastage in accordance with the XIAFLEX® Prescribing Information.

It is possible that any given payor may accept or require a different coding paradigm for same-day, dual cord treatments such as the use of modifier 51, 76, 59, or XS and/or billing service individually on separate line items. Please contact the XIAFLEX® Reimbursement Helpline (1-877-942-3539) or the payor's Provider Service Representative to acquire more information on coding guidance.

INDICATION

XIAFLEX® is indicated for the treatment of adult patients with Dupuytren's contracture with a palpable cord.

IMPORTANT SAFETY INFORMATION FOR XIAFLEX

- XIAFLEX is contraindicated in patients with a history of hypersensitivity to XIAFLEX or to collagenase used in any other therapeutic application or application method
- In the controlled and uncontrolled portions of clinical trials in Dupuytren's contracture, flexor tendon ruptures occurred after XIAFLEX injection. Injection of XIAFLEX into collagen-containing structures such as tendons or ligaments of the hand may result in damage to those structures and possible permanent injury such as tendon rupture or ligament damage. Therefore, XIAFLEX should be injected only into the collagen cord with a metacarpophalangeal (MP) or proximal interphalangeal (PIP) joint contracture, and care should be taken to avoid injecting into tendons, nerves, blood vessels, or other collagen-containing structures of the hand. When injecting a cord affecting a PIP joint of the fifth finger, the needle insertion should not be more than 2 to 3 mm in depth and avoid injecting more than 4 mm distal to the palmar digital crease
- Other XIAFLEX-associated serious local adverse reactions in the controlled and uncontrolled portions of the clinical studies included pulley rupture, ligament injury, complex regional pain syndrome (CRPS), sensory abnormality of the hand, and skin laceration (tear). In a historically controlled post-marketing trial, the incidence of skin laceration (22%) was higher for subjects treated with two concurrent injections of XIAFLEX compared with subjects treated with up to three single injections in the placebo-controlled premarketing trials (9%). Post-marketing cases of skin laceration requiring skin graft after finger extension procedures and local skin and soft-tissue necrosis, some requiring skin grafting, or other surgical interventions including finger amputation have been reported. Signs or symptoms that may reflect serious injury to the injected finger/hand should be promptly evaluated because surgical intervention may be required
- Cases of syncope and presyncope have been reported in the post-marketing period in patients treated with XIAFLEX. In most cases in patients with Dupuytren's contracture, the injection procedure, finger extension procedure, or pain following the procedures were reported as potential triggers for the events, suggesting a vasovagal mechanism. Most, but not all, cases occurred in the immediate treatment period (injection or finger extension procedure) or within 1 to 2 days following the injection or finger extension procedure. If presyncopal symptoms occur, patients should remain recumbent until symptoms resolve. Syncope may be associated with bodily injuries, including concussion, head abrasion, and other accidental injuries
- In the controlled portions of the clinical trials in Dupuytren's contracture, a greater proportion of XIAFLEX-treated patients (15%) compared to placebo-treated patients (1%) had mild allergic reactions (pruritus) after up to 3 injections. The incidence of XIAFLEX-associated pruritus increased after more XIAFLEX injections in patients with Dupuytren's contracture
- Because XIAFLEX contains foreign proteins, severe allergic reactions to XIAFLEX can occur. Anaphylaxis was reported in a post-marketing clinical trial in one patient who had previous exposure to XIAFLEX for the treatment of Dupuytren's contracture. Healthcare providers should be prepared to address severe allergic reactions following XIAFLEX injections
- In the XIAFLEX trials in Dupuytren's contracture, 70% and 38% of XIAFLEX-treated patients developed an ecchymosis/contusion or an injection site hemorrhage, respectively. Patients with abnormal coagulation (except for patients taking low-dose aspirin, eg, up to 150 mg per day) were excluded from participating in these studies. Therefore, the efficacy and safety of XIAFLEX in patients receiving anticoagulant medications (other than low-dose aspirin, eg, up to 150 mg per day) within 7 days prior to XIAFLEX administration is not known. In addition, it is recommended to avoid use of XIAFLEX in patients with coagulation disorders, including patients receiving concomitant anticoagulants (except for low-dose aspirin)
- In the XIAFLEX clinical trials for Dupuytren's contracture, the most common adverse reactions reported in $\geq 25\%$ of patients treated with XIAFLEX and at an incidence greater than placebo were edema peripheral (eg, swelling of the injected hand), contusion, injection site hemorrhage, injection site reaction, and pain in the injected extremity
- Post-marketing experience – Syncope and presyncope have been reported in patients treated with XIAFLEX. Most, but not all, cases occurred in the immediate treatment period or within 1 to 2 days following injection. Bodily injuries associated with the syncopal events have been reported

Click for full [Prescribing Information](#), including [Medication Guide](#).

XIAFLEX®
collagenase clostridium histolyticum

 **endo**
pharmaceuticals
an endo international company

XIAFLEX® is a registered trademark of Endo International plc or one of its affiliates.
© 2022 Endo International plc or one of its affiliates. All rights reserved.
MM-05791/August 2022 www.xiaflex.com 1-800-462-ENDO (3636)