XIAFLEX® (collagenase clostridium histolyticum)
Coding Scenario 2: Procedure Modifier Required

HEALTH INSURANCE CLAIM FORM

APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12

Medicare Part B Claims

CARRIER

PICA

1. MEDICARE (Medicare#)	MEDICAID (Medicaid#)	TRICARE (ID#/DoD#)	CHAMPVA (Member ID#)	GROUP HEALTH PLAN (ID#)	FECA BLK LUNG (ID#)	OTHER (ID#)

1a. INSURED'S I.D. NUMBER (For Program in Item 1): 123-45-6789A

2. PATIENT'S NAME (Last Name, First Name, Middle Initial)	3. PATIENT'S BIRTH DATE MM DD YY / SEX	4. INSURED'S NAME (Last Name, First Name, Middle Initial)
Doe, John Q.	10 19 1935 / M X F	Same
5. PATIENT'S ADDRESS (No., Street)	6. PATIENT RELATIONSHIP TO INSURED	7. INSURED'S ADDRESS (No., Street)
1212 Main St.	Self X Spouse Child Other	Same
CITY: Any City / STATE: XX	8. RESERVED FOR NUCC USE	CITY / STATE
ZIP CODE: XXXXX / TELEPHONE (Include Area Code)		ZIP CODE / TELEPHONE (Include Area Code)

PATIENT AND INSURED INFORMATION

9. OTHER INS

a. OTHER INS: AARP

b. RESERVED

c. RESERVED

d. INSURANC

12. PATIENT'S to process below. SIGNED

d 9d.

uthorize supplier for

14. DATE OF MM DD

PATION YY

17. NAME OF

17b. NPI

ICES YY

FROM TO

19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)

20. OUTSIDE LAB? YES NO $ CHARGES

21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Relate A-L to service line below (24E) ICD Ind.

A. M72.0 B. C. D.

E. F. G. H.

I. J. K. L.

22. RESUBMISSION CODE ORIGINAL REF. NO.

23. PRIOR AUTHORIZATION NUMBER

24. A. DATE(S) OF SERVICE From MM	DD	YY	To MM	DD	YY	B. PLACE OF SERVICE	C. EMG	D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances) CPT/HCPCS	MODIFIER	E. DIAGNOSIS POINTER	F. $ CHARGES	G. DAYS OR UNITS	H. EPSDT Family Plan	I. ID. QUAL.	J. RENDERING PROVIDER ID. #
1	10	03	21	10	03	21	11		J0775		A	XXX XX	116		NPI	
2	10	03	21	10	03	21	11		J0775	JW	A	XXX XX	64		NPI	
3	10	03	21	10	03	21	11		20527		A	XXX XX	1		NPI	
4	10	03	21	10	03	21	11		20527		A	XXX XX	1		NPI	
5	10	05	21	10	05	21	11		26341		A	XXX XX	1		NPI	
6	10	05	21	10	05	21	11		26341		A	XXX XX	1		NPI	

Use JW modifier to indicate how much drug is wasted from single-use vials.

These amounts represent 58 units per cord/joint (total of 116 units) and wastage of 32 units per cord/joint (total of 64 units) and are provided as an example of the recommended dose and wastage in accordance with the XIAFLEX® Prescribing Information.

25. FEDERAL TAX I.D. NUMBER SSN EIN	26. PATIENT'S ACCOUNT NO.	27. ACCEPT ASSIGNMENT? (For govt. claims, see back)	28. TOTAL CHARGE	29. AMOUNT PAID	30. Rsvd for NUCC Use
123456789		YES NO	$ XXXX XX	$	XXX XX

31. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREES OR CREDENTIALS (I certify that the statements on the reverse apply to this bill and are made a part thereof.)

SIGNED DATE

32. SERVICE FACILITY LOCATION INFORMATION

a. NPI b.

It is possible that any given payor may accept or require a different coding paradigm for same-day, dual cord treatments such as the use of modifier 51, 76, 59, or XS and/or billing service individually on separate line items. Please contact the XIAFLEX® Reimbursement Helpline (1-877-942-3539) or the payor's Provider Service Representative to acquire more information on coding guidance.

NUCC Instruction Manual available at: www.nucc.org PLEASE PRINT OR ...0 (02-12)

SAMPLE 1500 FORM
TREATMENT BILLING
WHEN TREATING 2 CORDS/JOINTS

Based on prior policy, this sample represents how your Medicare Administrative Contractor (MAC) is likely to require completion of claim forms for J0775, and the CPT® codes 20527 and 26341. This sample form is not intended to replace or modify your MAC's policy, and use of this form does not guarantee payment or take the place of professional coding advice. Coding is part of the clinical decision, and each provider is responsible for selecting the billing codes that most accurately describe the services provided, and for adhering to all payor guidance. Information is subject to change. Please refer to the policy on the MAC's website. This sample claim form does not represent any clinical or treatment recommendation.

Please see Indication and Important Safety Information on next page.

Click for full Prescribing Information and Medication Guide.

INDICATION

XIAFLEX® is indicated for the treatment of adult patients with Dupuytren's contracture with a palpable cord.

IMPORTANT SAFETY INFORMATION FOR XIAFLEX

- XIAFLEX is contraindicated in patients with a history of hypersensitivity to XIAFLEX or to collagenase used in any other therapeutic application or application method
- In the controlled and uncontrolled portions of clinical trials in Dupuytren's contracture, flexor tendon ruptures occurred after XIAFLEX injection. Injection of XIAFLEX into collagen-containing structures such as tendons or ligaments of the hand may result in damage to those structures and possible permanent injury such as tendon rupture or ligament damage. Therefore, XIAFLEX should be injected only into the collagen cord with a metacarpophalangeal (MP) or proximal interphalangeal (PIP) joint contracture, and care should be taken to avoid injecting into tendons, nerves, blood vessels, or other collagen-containing structures of the hand. When injecting a cord affecting a PIP joint of the fifth finger, the needle insertion should not be more than 2 to 3 mm in depth and avoid injecting more than 4 mm distal to the palmar digital crease
- Other XIAFLEX-associated serious local adverse reactions in the controlled and uncontrolled portions of the clinical studies included pulley rupture, ligament injury, complex regional pain syndrome (CRPS), sensory abnormality of the hand, and skin laceration (tear). In a historically controlled post-marketing trial, the incidence of skin laceration (22%) was higher for subjects treated with two concurrent injections of XIAFLEX compared with subjects treated with up to three single injections in the placebo-controlled premarketing trials (9%). Post-marketing cases of skin laceration requiring skin graft after finger extension procedures and local skin and soft-tissue necrosis, some requiring skin grafting, or other surgical interventions including finger amputation have been reported. Signs or symptoms that may reflect serious injury to the injected finger/hand should be promptly evaluated because surgical intervention may be required
- In the controlled portions of the clinical trials in Dupuytren's contracture, a greater proportion of XIAFLEX-treated patients (15%) compared to placebo-treated patients (1%) had mild allergic reactions (pruritus) after up to 3 injections. The incidence of XIAFLEX-associated pruritus increased after more XIAFLEX injections in patients with Dupuytren's contracture
- Because XIAFLEX contains foreign proteins, severe allergic reactions to XIAFLEX can occur. Anaphylaxis was reported in a post-marketing clinical trial in one patient who had previous exposure to XIAFLEX for the treatment of Dupuytren's contracture. Healthcare providers should be prepared to address severe allergic reactions following XIAFLEX injections
- In the XIAFLEX trials in Dupuytren's contracture, 70% and 38% of XIAFLEX-treated patients developed an ecchymosis/contusion or an injection site hemorrhage, respectively. Patients with abnormal coagulation (except for patients taking low-dose aspirin, eg, up to 150 mg per day) were excluded from participating in these studies. Therefore, the efficacy and safety of XIAFLEX in patients receiving anticoagulant medications (other than low-dose aspirin, eg, up to 150 mg per day) within 7 days prior to XIAFLEX administration is not known. In addition, it is recommended to avoid use of XIAFLEX in patients with coagulation disorders, including patients receiving concomitant anticoagulants (except for low-dose aspirin)
- In the XIAFLEX clinical trials for Dupuytren's contracture, the most common adverse reactions reported in ≥25% of patients treated with XIAFLEX and at an incidence greater than placebo were edema peripheral (eg, swelling of the injected hand), contusion, injection site hemorrhage, injection site reaction, and pain in the injected extremity. Post-marketing experience – Syncope and presyncope have been reported in patients treated with XIAFLEX for Dupuytren's contracture. In some cases, pain from injection and pain during finger extension procedures were identified as potential triggers for syncopal events

Click for full Prescribing Information and Medication Guide.

XIAFLEX®
collagenase clostridium histolyticum

endo pharmaceuticals
an endo international company

MM-05677/January 2022 www.xiaflex.com 1-800-462-ENDO (3636)