

1 Hometown Hospital 123 Main Street Anycity, Anystate 12345		2	3a PAT. CNTL. # b. MED. REC. #	4 TYPE OF BILL
8 PATIENT NAME a Doe, John Q.		9 PATIENT ADDRESS a Anycity, Anystate 12345		
10 BIRTHDATE	11 SEX	12 DATE	ADMISSION 13 HR 14 TYPE 15 SRC	16 DHR
31 OCCURRENCE CODE		32 OCCURRENCE DATE	33 OCCURRENCE CODE	34 OCCURRENCE DATE
35 OCCURRENCE SPAN FROM		36 OCCURRENCE SPAN THROUGH		37
38		39 VALUE CODES AMOUNT	40 VALUE CODES AMOUNT	41 VALUE CODES AMOUNT
42 REV. CD.	43 DESCRIPTION	44 HCPCS / RATE / HIPPS CODE	45 SERV. DATE	46 SERV. UNITS
1 0250	Drugs and Biologicals	J0775	012016	90
2 0342	Therapeutic Procedure	20527	012016	1
3 0342	Therapeutic Procedure	26341	012016	1
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				
19				
20				
21				
22				
23				
50 PAYER N				
58 INSURED'S NAME	59 PREL	60 INSURED'S UNIQUE ID	61 GROUP NAME	62 INSURANCE GROUP NO.
63 TREATMENT AUTHORIZATION CODES	64 DOCUMENT CONTROL NUMBER		65 EMPLOYER NAME	
66 DX	M72.0	68		
69 ADMIT DX	70 PATIENT REASON DX	71 PPS CODE	72 ECI	73
74 PRINCIPAL PROCEDURE CODE	a. OTHER PROCEDURE CODE	b. OTHER PROCEDURE CODE	75	76 ATTENDING NPI
				QUAL
				LAST
				FIRST
				77 OPERATING NPI
				QUAL
				LAST
				FIRST
80 REMARKS	81 CC a		78 OTHER NPI	QUAL
	b		LAST	FIRST
	c		79 OTHER NPI	QUAL
	d		LAST	FIRST

**SAMPLE 1450 FORM  
SINGLE CORD TREATMENT**

Based on prior policy, this sample represents how your Medicare Administrative Contractor (MAC) is likely to require completion of claim forms for J0775, and the CPT® codes 20527 and 26341. This sample form is not intended to replace or modify your MAC's policy, and use of this form does not guarantee payment or take the place of professional coding advice. Coding is part of the clinical decision, and each provider is responsible for selecting the billing codes that most accurately describe the services provided and for adhering to all payor guidance. Information is subject to change. Please refer to the policy on the MAC's Web site. This sample claim form does not represent any clinical or treatment recommendation.

## INDICATION

XIAFLEX® (collagenase clostridium histolyticum) is indicated for the treatment of adult patients with Dupuytren's contracture with a palpable cord.

## IMPORTANT SAFETY INFORMATION FOR XIAFLEX®

- XIAFLEX® is contraindicated in patients with a history of hypersensitivity to XIAFLEX® or to collagenase used in any other therapeutic application or application method
- In the controlled and uncontrolled portions of clinical trials in Dupuytren's contracture, flexor tendon ruptures occurred after XIAFLEX® injection. Injection of XIAFLEX® into collagen-containing structures such as tendons or ligaments of the hand may result in damage to those structures and possible permanent injury such as tendon rupture or ligament damage. Therefore, XIAFLEX® should be injected only into the collagen cord with a MP or PIP joint contracture, and care should be taken to avoid injecting into tendons, nerves, blood vessels, or other collagen-containing structures of the hand. When injecting a cord affecting a PIP joint of the fifth finger, the needle insertion should not be more than 2 to 3 mm in depth and avoid injecting more than 4 mm distal to the palmar digital crease
- Other XIAFLEX®-associated serious local adverse reactions in the controlled and uncontrolled portions of the studies included pulley rupture, ligament injury, complex regional pain syndrome (CRPS), sensory abnormality of the hand, and skin laceration (tear). In a historically controlled post-marketing trial, the incidence of skin laceration (22%) was higher for subjects treated with two concurrent injections of XIAFLEX® compared with subjects treated with up to three single injections in the placebo-controlled premarketing trials (9%). Cases of skin laceration requiring skin graft after finger extension procedures have been reported post-marketing. Signs or symptoms that may reflect serious injury to the injected finger/hand should be promptly evaluated because surgical intervention may be required
- In the controlled portions of the clinical trials in Dupuytren's contracture, a greater proportion of XIAFLEX®-treated patients (15%) compared to placebo-treated patients (1%) had mild allergic reactions (pruritus) after up to 3 injections. The incidence of XIAFLEX®-associated pruritus increased after more XIAFLEX® injections in patients with Dupuytren's contracture
- Because XIAFLEX® contains foreign proteins, severe allergic reactions to XIAFLEX® can occur. Anaphylaxis was reported in a post-marketing clinical study in one patient who had previous exposure to XIAFLEX® for the treatment of Dupuytren's contracture. Healthcare providers should be prepared to address severe allergic reactions following XIAFLEX® injections
- In the XIAFLEX® trials in Dupuytren's contracture, 70% and 38% of XIAFLEX®-treated patients developed an ecchymosis/contusion or an injection site hemorrhage, respectively. Patients with abnormal coagulation (except for patients taking low-dose aspirin, eg, up to 150 mg per day) were excluded from participating in these studies. Therefore, the efficacy and safety of XIAFLEX® in patients receiving anticoagulant medications (other than low-dose aspirin, eg, up to 150 mg per day) within 7 days prior to XIAFLEX® administration is not known. In addition, it is recommended to avoid use of XIAFLEX® in patients with coagulation disorders, including patients receiving concomitant anticoagulants (except for low-dose aspirin)
- In the XIAFLEX® clinical trials for Dupuytren's contracture, the most common adverse reactions reported in ≥25% of patients treated with XIAFLEX® and at an incidence greater than placebo were edema peripheral (eg, swelling of the injected hand), contusion, injection site hemorrhage, injection site reaction, and pain in the injected extremity

Please [click here](#) for full Prescribing Information, including Medication Guide.

**XIAFLEX®**  
collagenase clostridium histolyticum

 **endo**  
pharmaceuticals  
an endo international company

**Rx Only**

XIAFLEX® is a registered trademark of Endo International plc or one of its affiliates.  
© 2016 Endo Pharmaceuticals Inc. All rights reserved. Malvern, PA 19355  
**XD-03534(1)/January 2016** www.xiaflex.com 1-800-462-ENDO (3636)